

Hypoglycemia Symptoms

Circle all that Apply

Mild Range _____	Moderate Range _____	Severe Range _____
Hunger, Pallor, Dizziness Irritability, Crying, Shakiness, Anxiety Drowsiness, Headache	Sleepiness, Erratic Behavior Confusion, Slurred Speech Poor Coordination	Pain/Discomfort with Swallowing Combativeness, Unconsciousness Seizures
Treatment		
<ol style="list-style-type: none"> 1. Provide sugar source: 3-4 glucose tabs, OR 4 oz juice, OR 4 oz regular soda, OR glucose gel (15g) 2. Wait 10-15 minutes 3. Retest glucose, if less than ____ retreat with sugar source every 15 minutes until glucose over _____ 4. Student may return to class when BSL is over _____ 	<ol style="list-style-type: none"> 1. Provide sugar source: 3-4 glucose tabs, OR 4 oz juice, OR 4 oz regular soda, OR glucose gel (15g) 2. Wait 10-15 minutes 3. Retest glucose, if less than ____ retreat with sugar source every 15 minutes until glucose over _____ 4. Notify parent 	<ol style="list-style-type: none"> 1. <u>CALL 911</u> 2. Give glucagon as ordered 3. Position on side 4. Contact parent

Hyperglycemia Symptoms

Circle all that Apply

Early Signs	Later Signs	Call 911
Frequent urination, Increased thirst Blurred Vision, Fatigue, Headache	Fruity-smelling breath, Weakness Nausea/Vomiting, Shortness of Breath, Dry mouth, Confusion Coma, Abdominal Pain	Inability to hold down any food or fluids BSL persistently above 240 and ketones are present in urine